

学校编码：10384

学号：20620100153627

厦门大学

博士学位论文

聚苯乙烯负载型和锌-腺嘌呤骨架型材料的构建及其催化性能研究

Construction of Polystyrene-supported and  
Zinc-Adeninate Framework Materials and  
Investigation of Their Catalytic  
Performance

王结祥

指导教师：陈秉辉

专业名称：工业催化

答辩日期：2013年9月

## 厦门大学学位论文原创性声明

本人呈交的学位论文是本人在导师指导下，独立完成的研究成果。本人在论文写作中参考其他个人或集体已经发表的研究成果，均在文中以适当方式明确标明，并符合法律规范和《厦门大学研究生学术活动规范(试行)》。

另外，该学位论文为( )课题(组)的研究成果，获得( )课题(组)经费或实验室的资助，在( )实验室完成。(请在以上括号内填写课题或课题组负责人或实验室名称，未有此项声明内容的，可以不作特别声明。)

声明人(签名)：

年 月 日

# 厦门大学学位论文著作权使用声明

本人同意厦门大学根据《中华人民共和国学位条例暂行实施办法》等规定保留和使用此学位论文，并向主管部门或其指定机构送交学位论文(包括纸质版和电子版)，允许学位论文进入厦门大学图书馆及其数据库被查阅、借阅。本人同意厦门大学将学位论文加入全国博士、硕士学位论文共建单位数据库进行检索，将学位论文的标题和摘要汇编出版，采用影印、缩印或者其它方式合理复制学位论文。

本学位论文属于：

(        )1. 经厦门大学保密委员会审查核定的保密学位论文，于  
年 月 日解密，解密后适用上述授权。

(        )2. 不保密，适用上述授权。

(请在以上相应括号内打“√”或填上相应内容。保密学位论文应是已经厦门大学保密委员会审定过的学位论文，未经厦门大学保密委员会审定的学位论文均为公开学位论文。此声明栏不填写的，默认为公开学位论文，均适用上述授权。)

声明人(签名)：

年    月    日

## 摘 要

在倡导可持续发展和绿色能源化学背景下，工业催化朝着少分离、低能耗的多相催化发展，同时也向高附加值的精细化工发展。特别是面对石化枯竭和能源结构过于单一化的隐忧，催化一方面积极开拓合成新能源化学品的反应，如生物质转化等，另一方面也致力于从碳一化学的角度拓展碳链增长的反应丰富合成路线，为可能的能源危机做技术储备。其中，CO<sub>2</sub>作为丰富安全的碳资源，在温室效应等环境问题日益恶化的情况下更加受关注，对其合理地催化转化将有助实现全球碳循环并丰富能源结构。

本文以工业应用较成熟的聚苯乙烯型离子交换树脂为起点，以低选择性的甲醛丙酮羟醛缩合制较不稳定化合物丁酮醇的碳链增长碱催化反应为探针入手。尝试表面活性剂的引入和设计索氏提取反应，发现提高反应速率和实现中间目标产物丁酮醇和催化剂的分离，都能有效避免深度反应的进行，产率得以提高。进一步地，对商业化树脂进行改性，主要通过以下途径：（1）对叔胺树脂通过Menshutkin氮烷基化反应在叔胺氮上引入长碳链；（2）在叔胺基团存在下，对聚苯乙烯骨架苯环上通过Friedel-Crafts烷基化反应引入长链酰基，并接着通过Menshutkin反应在叔胺上引入长碳链，从而构成双长碳链。催化结果表明，通过途径（1），丁酮醇产率由44.3%上升超过50%；而通过途径（2），产率达到71.3%。气质分析显示，弱碱叔胺相对于强碱季铵树脂，其催化副产物减少，而改良后的树脂其催化副产物进一步下降。CO<sub>2</sub>-TPD和HCHO-TPD表明，随着吸附分子尺寸的增大，在改性树脂上的吸附量明显下降，这预示着从高分子微观角度而言，长链的引入有效提高高分子内部位阻效应，使尺寸较大的丁酮醇更易从活性位点脱附下来，从而提高其选择性。

然而，上述接枝离子交换树脂受胺基影响，其热稳定性较差。我们引入含氮杂环负载于聚苯乙烯上，有效提高其热稳定性达近400℃。该材料在CO<sub>2</sub>合成环状碳酸酯上是一种高效的多相催化材料。烷基胺和卤代锌的引入均有利于活性的提高，但热稳定性因此有所下降。考察不同氮杂环对活性的影响，五元氮杂环的活性明显高于六元环。进而，通过衰减全反射傅里叶变换红外和核磁共振分析，考察产物碳酸丙烯酯（PC）和催化活性基团咪唑环（Im）的相互作用，从反应终态间接地反向推导

催化过程，提出一种可行的反应机理。碳酸丙烯酯和咪唑环均是平面五元环结构，从电子诱导效应而言，PC上的O-C-O结构正好与Im上含孤对电子的N-C=N配对。并进一步推测，在咪唑环上实现了CO<sub>2</sub>和环氧丙烷（PO）的吸附、以及PO开环和CO<sub>2</sub>的活化插入过程，相当于两个分子在咪唑环上的催化组装过程。催化结果表明，转化率达99.7%，选择性99.6%，TON达204以上。并且重复利用四次未发现活性有明显下降。

进一步地，高聚物负载量相对有限，对大宗CO<sub>2</sub>转化略显吃力，于是我们引入具有高比表面、富氮的金属有机骨架材料锌-腺嘌呤-异烟酸（Zn-Ad-Int）的构建。尝试晶体生长温度、锌盐中阴离子变化、阳离子镧掺杂、烷基胺接枝和KI添加等手段，了解到亲核试剂对PO的进攻、开环和CO<sub>2</sub>的插入加成影响重大。从而引入ZnX<sub>2</sub>-Ad-Int的构建，卤素对反应活性的影响依次有Cl<Br<I<sup>-</sup>，这在于碘离子既有较好的亲核性又有很好的离去性。该材料耐热性达280℃，重复利用后活性保持在96.7%左右。值得一提的是，水介质生长的ZnX<sub>2</sub>-Ad-Int-H<sub>2</sub>O热稳定性高达近400℃。

前面，我们更多的是着眼于催化材料，而最后一章则从反应入手。为了拓展CO<sub>2</sub>的催化转化应用，我们从CO<sub>2</sub>和甲醇直接合成碳酸二甲酯（DMC）研究起，通过引入疏水性的骨架材料——聚苯乙烯和金属有机骨架（MOFs），一定程度上促进了DMC的生成。相比于Zn-W-Mim，疏水性骨架Zn(bdc)(ted)<sub>0.5</sub>对铜基催化剂的负载更有利于DMC的生成。另外，在高温高压酸碱环境下，骨架材料会有一定程度的降解。苯环的活化生成苯甲酸甲酯和水等也进一步抑制了竞争反应DMC的生成。而聚苯乙烯的降解又为该反应注入了新的意义。另一方面，通过对CO<sub>2</sub>转化中较难实现反应的摸索，我们有机会将聚苯乙烯负载和金属有机骨架材料两者的一些改性手段应用其中，如不同配体的负载、金属螯合、疏水性MOFs的应用和杂化型MOFs的尝试等等。

**关键词：**固体碱；离子交换树脂；聚苯乙烯；羟醛缩合；丁酮醇；氮杂环；二氧化碳；碳酸丙烯酯；腺嘌呤；金属有机骨架；碳酸二甲酯

## Abstract

Under the wake of the sustainable development and green energy chemistry, industrial catalysis parades towards the heterogeneous catalysis with less separation and low energy consumption, as well as the fine chemical industry with high additional value. To relieve the hidden threat and panics of exhausting fossil energy and the limited energy options, researches on catalysis concentrate efforts on the reactions exploring for new energy chemicals such as biomass conversion, as well as on C1 chemistry to expand carbon chain elongation reactions for riching synthetic routes, targeting to have preparations for the potential energy crisis. Thereinto, as a rich and safe carbon resource, CO<sub>2</sub> attracts people's attentions for climatic issues such as the greenhouse effect. CO<sub>2</sub> catalytic conversion constitutes a key link of the global carbon cycle, and can also improve the energy structure.

This work begins with a commercial polystyrene type ion exchange resin, and takes the aldol condensation of formaldehyde and acetone as the probe reaction, which is a carbon elongation reaction with less stable products and low selectivity. Surfactant and Soxhlet extraction are associated to improve the yield. It is inferred that both the improvement of reaction rate and the separation of the medium product 4-hydroxy-2-butanone from catalysts could reduce the side reactions, thus increased the yield. Subsequently, two types of modifications for resins are carried out: (1) one long alkyl chain on the copolymer tertiary amine via Menshutkin N-alkylation reaction; (2) two long alkyl chains: one on polystyrene aromatic ring through Friedel–Crafts alkylation under the circumstance of tertiary amine and the other on the copolymer tertiary amine using the Menshutkin N-alkylation reaction. The reaction results indicate that the yield for 4-hydroxy-2-butanone can be increased from 44.3% to around 50% for the resin with one long alkyl chain (the first case), while this can achieve up to 71.3% for the resin with

two long alkyl chains. The GC-MS analysis shows that the weaker alkali tertiary amine resin has fewer by-products than stronger alkali quaternary amine resin, and the content of by-products of modified resin decreases sharply. CO<sub>2</sub>-TPD and HCHO-TPD prove that the adsorbing capacities on the modified resins are decreasing with the size of the adsorbed molecule increasing. It indicates that adding long chain effectively improves polymer internal steric effect in the view of the polymer microstructure, then the bigger molecular 4-hydroxy-2-butanone is more likely to desorb from the active sites, thus the selectivity is improved.

However, the thermal stability of ion exchange resins is weakening by adding amine groups. N-heterocycles are introduced to load on polystyrene and the thermal stability can reach approximately 400 °C with such grafting. They are proved to be an efficient heterogeneous catalyst for converting CO<sub>2</sub> into cyclic carbonate. The introduction of alkyl amines and zinc halides can both boost the activity, but it decreases the thermostability of the catalysts. A series of N-heterocycles are chose to catalyze CO<sub>2</sub> cycloaddition. Five-membered heterocyclic compounds exhibit higher activities than six-membered ones. The interaction between the product propylene carbonate (PC) and the catalytic active group imidazole ring (Im) is studied through ATR-IR and NMR. Then, the catalytic process is inferred backward from the final state of the reaction. PC and Im are both perfectly planar with a five-membered ring structure. Based on the electron induced effect, the O-C-O of PC can match with the N-C=N of imidazole, which owns unpaired nitrogens. A plausible mechanism is proposed. CO<sub>2</sub> and PO are adsorbed on an imidazole ring, then subsequently with the ring-opening of PO and the activation and insertion of CO<sub>2</sub> into PO, just like a catalytic assembly process of two moleculars on one imidazole ring. Finally, the yield reaches 99.7%, with the selectivity 99.6% and TON above 204. The catalyst can be reused over four times without appreciable loss of catalytic activity.

Further, bulk CO<sub>2</sub> conversion seems to be slightly tired due to the low loading of

polymer-supported materials and the Metal-Organic Frameworks (MOFs) may overcome such difficulty based on the current advantages. As a part of this work, a Zn-Ad-Int framework is constructed with high surface area and nitrogen-rich. The effects of MOF components on CO<sub>2</sub> conversion are studied, including growth temperature, different zinc salts, Lanthanum embedded, alkyl amine grafting and KI addition. We find that the attack of nucleophilic reagent on PO is a key point for PO ring-opening and CO<sub>2</sub> inserting. Then, we focus on the construction of ZnX<sub>2</sub>-Ad-Int (X=Cl, Br, I). The activity increases as followed: Cl<Br<I-, which is because I- owns a good nucleophilic ability, as well as a good leaving ability. Their thermostability can reach 280, and the recycling activity can keep at about 96.7%. It is worth to mention that the thermal stability of ZnX<sub>2</sub>-Ad-Int-H<sub>2</sub>O can reach approximately 400.

Prior to this, we shift our attention onto the reaction in the last chapter, as focusing on catalytic materials previously. For expanding CO<sub>2</sub> catalytic application, we start our new research target on the direct synthesis of DMC from CO<sub>2</sub> and methanol. The introduction of hydrophobic frameworks, such as polystyrene and MOFs, could promote DMC yield. Compared with Zn-W-Mim, Cu embeds into hydrophobic Zn(bdc)(ted)<sub>0.5</sub> framework is more conducive to yield DMC. Moreover, under the acid and alkali circumstance with high temperature and high pressure, both the two frameworks, polystyrene and Zn(bdc)(ted)<sub>0.5</sub> suffer with degradation. The activation of benzene ring to produce methyl benzoate and water may further suppress DMC yield. On the other hand, it also brings a new significance on the reclamation of waste polystyrene. In addition, during the exploration for difficult CO<sub>2</sub> conversion reactions, we practice a series of modification methods on polystyrene-supported materials and MOFs, such as different ligands loading, metal chelation, hydrophobic MOF and hybrid framework applications.

**Keywords:** solid base; ion exchange resin; polystyrene; aldol condensation; 4-



hydro-2-butanone; N-heterocycle; carbon dioxide; propylene carbonate; adenine;  
metal-organic framework (MOF); dimethyl carbonate.

厦门大学博硕士论文摘要库

## 参考资料

- [1]. Haag W.O. and Hill C. Sulfonated resin catalyst with enhanced activity[P]. US3326866, Belgian, 1967: 686-721.
- [2]. Morimoto H., Yukawa T., Lu G., et al. Lewis base assisted Bronsted base catalysis: bidentate phosphine oxides as activators and modulators of Bronsted basic lanthanum-aryloxides[J]. *Angewandte Chemie International Edition*, 2008. 47(47): 9125-9129.
- [3]. 唐新硕, 王新平. 催化科学发展及其理论[M]. 杭州: 浙江大学出版社. 2012.
- [4]. Anastas P T and Warner J. C. Green Chemistry: Theory and Practice [M]. Oxford: Oxford University Press. 1998.
- [5]. Wight A P and Davis M E. Design and Preparation of Organic-Inorganic Hybrid Catalysts[J]. *Chemical Reviews*, 2002. 102(10): 3589-3614.
- [6]. Clapham B, Reger T. S, and Janda K. D. Polymer-supported catalysis in synthetic organic chemistry[J]. *Tetrahedron*, 2001. 57: 4637-4662.
- [7]. Benaglia M., and Cozzi F.. Polymer-Supported Organic Catalysts[J]. *Chemical Reviews*, 2003. 103(9): 3401-3430.
- [8]. Buchmeiser M. R. polymer-supported well-defined.[J]. *Chemical Reviews*, 2009. 109(2): 303-321.
- [9]. Haraguchi M, Nakagawa T, and Nose T. Miscibility of associated polymer blend solutions: 1. One-end-aminated polystyrene-one-end-carboxylated poly(ethylene glycol) blends in toluene[J]. *Polymer*, 1995. 36(13): 2567-2572.
- [10]. 钱延龙, 陈新滋. 金属有机化学与催化[M]. 北京: 化学工业出版社. 1997.
- [11]. 王结祥, 王强, 李云华等. 聚苯乙烯树脂催化羟醛缩合研究进展[J]. *化工进展*, 2011. 30(7): 1521-1528.
- [12]. Denmark S E and Beutner G L. Lewis base catalysis in organic synthesis[J]. *Angewandte Chemie International Edition*, 2008. 47(9): 1560-1638.
- [13]. Figueras F. Base catalysis in the synthesis of fine chemicals[J]. *Topics in Catalysis*, 2004. 29(3-4): 189-196.
- [14]. Figueras F, Kantam M. L., and Choudary B. M. Solid Base Catalysts in Organic Synthesis[J]. *Current Organic Chemistry*, 2006. 10(13): 1627-1637.
- [15]. Leyva C, Trejo F., and Ancheyta J., et al. On the Use of Acid-Base-Supported Catalysts for Hydroprocessing of Heavy Petroleum[J]. *Industrial & Engineering Chemistry Research*, 2007. 46: 7448-7466.
- [16]. Ono Y. Solid base catalysts for the synthesis of fine chemicals[J]. *Journal of Catalysis*, 2003. 216(1-2): 406-415.
- [17]. Hattori H. Solid base catalysts- generation of basic sites and application to organic synthesis[J]. *Applied Catalysis A: General*, 2001. 222: 247-259.
- [18]. Hattori H. solid base catalysts: generation, characterization, and catalytic behavior of basic sites[J]. *Journal of the Japan Petroleum Institute*, 2004. 47(2): 67-81.
- [19]. Tanabe K and Holderich W. F. Industrial application of solid acid-base catalysts[J]. *Applied Catalysis A: General*, 1999. 181(2): 399-434.
- [20]. Weitkamp J, Hunger M., and Rymas U. Base catalysis on microporous and mesoporous materials-recent progress and perspectives[J]. *Microporous and Mesoporous Materials*, 2001. 48: 255-270.
- [21]. Davis R J. New perspectives on basic zeolites as catalysts and catalyst supports[J]. *Journal of Catalysis*, 2003. 216(1-2): 396-405.
- [22]. 梁泽磊, 伍艳辉. 羟醛缩合体系反应精馏研究进展[J]. *化工时刊*, 2010. 24(1): 66 - 70.
- [23]. 蔡红, 周斌. 离子交换树脂在有机催化反应中的应用进展[J]. *化工进展*, 2007. 26(3): 386-391.
- [24]. Abrams I and Mullar J. A history of the origin and development of macroporous ion-exchange resins[J]. *Reactive and Functional Polymers*, 1997. 35(1): 7-22.
- [25]. Farrall J., FrBchet. B. Two Important Steps in the Functionalization of Polystyrene Resins[J]. *Journal of Organic Chemistry*, 1976. 41(24): 3877.

- [26].Frechet. Application of Phase-Transfer Catalysis to the Chemical Modification of cross linked polystyrene resins[J]. *Journal of Organic Chemistry*, 1979. 44(11): 1774.
- [27].Ford W T. Mechanisms of Polymer-Supported Catalysis. 3. Ion Exchange limitations and macroporous polystyrene supports[J]. *Macromolecules*, 1982. 15: 1246.
- [28].Kirschning A, Monenschein H., and Wittenberg R. Functionalized Polymers-Emerging Versatile Tools for Solution-Phase Chemistry and Automated Parallel Synthesis[J]. *Angew. Chem. Int. Ed.*, 2001. 40: 650- 679.
- [29].Lu J and Toy P H. Organic Polymer Supports for Synthesis and for Reagent and Catalyst Immobilization[J]. *Chemical Reviews*, 2009. 109(2): 815 – 838.
- [30].Campelo J. M., Luque R., Marinas J. M., et al. Sustainable preparation of supported metal nanoparticles and their applications in catalysis[J]. *ChemSusChem*, 2009. 2(1): 18-45.
- [31].Cop é ret C and Basset J. M. Strategies to Immobilize Well-Defined Olefin Metathesis Catalysts: Supported Homogeneous Catalysis vs. Surface Organometallic Chemistry[J]. *Advanced Synthesis & Catalysis*, 2007. 349(1-2): 78-92.
- [32].Leadbeater E. N. and Marco M. Preparation of Polymer-Supported Ligands and Metal Complexes for Use in catalysis[J]. *Chemical Reviews*, 2002. 102(10): 3217-3274.
- [33].Madhavan N., Jones C.W., and Marcusweck. Rational Approach to Polymer-Supported Catalysts-Synergy between Catalytic Reaction Mechanism and Polymer Design[J]. *Accounts of chemical research*, 2008. 41(9): 1153-1165.
- [34].Corma A and Garcia H. Silica-Bound Homogenous Catalysts as Recoverable and Reusable Catalysts in Organic Synthesis[J]. *Advanced Synthesis & Catalysis*, 2006. 348(12-13): 1391-1412.
- [35].Lee C. E. Supported Chiral Catalysts on Inorganic Materials [J]. *Chemical Reviews*, 2002. 102(10): 3495-3524.
- [36].赵刚等. 绿色有机催化[M]. 绿色化学化工丛书. 中国石化出版社. 2005.
- [37].Climent M J, Garcia H., Lopez R., et al. Acid – Base Bifunctional Catalysts for the Preparation of Fine Chemicals: Synthesis of Jasminaldehyde[J]. *Journal of Catalysis*, 2001. 197(2): 385-393.
- [38].Paull D H, Scerba M.T., Danforth E.A., et al. Bifunctional Asymmetric Catalysis-Cooperative Lewis Acid-Base Systems[J]. *Accounts of Chemical Research*, 2008. 41(5): 655-663.
- [39].Higgins J. S., Lipson J. E. G., and White R. P. A simple approach to polymer mixture miscibility[J]. *Philos Trans A Math Phys Eng Sci*, 2010. 368(1914): 1009-1025.
- [40].Janda P. W. and Janda K. D. Recent advances in soluble polymer-supported catalysts, reagents and synthesis[J]. *Chemical Reviews*, 1999: 1917-1924.
- [41].Gravert D. J. and Janda K. D. Organic Synthesis on Soluble Polymer Supports Liquid-Phase Methodologies[J]. *Chemical Reviews*, 1997. 97(2): 489-509.
- [42].Corain B., Zecca M., and Jerabek K.. Catalysis and polymer networks — the role of morphology and molecular[J]. *Journal of Molecular Catalysis A: Chemical*, 2001. 177: 3-20.
- [43].Wang Z., Chen G., and Ding K. Self-Supported Catalysts[J]. *Chemical Reviews*, 2009. 109(2): 322-359.
- [44].Kralik M and Biffis A. Catalysis by metal nanoparticles supported on functional organic polymers[J]. *Journal of Molecular Catalysis A: Chemical*, 2001. 177: 113-138.
- [45].Liu Z, et al. Organization of Nanoparticles on Soft[J]. *Nano Letters*, 2002. 2(3): 219-224.
- [46].Sommer W. J. and Weck M. Supported N-heterocyclic carbene complexes in catalysis[J]. *Coordination Chemistry Reviews*, 2007. 251(5-6): 860-873.
- [47].Zinck P., et al. Functionalization of syndiotactic polystyrene[J]. *Progress in Polymer Science*, 2009. 34(4): 369-392.
- [48].Kei Manabe, et al. Dehydrative Esterification of Carboxylic Acids with Alcohols Catalyzed by Polymer-Supported Sulfonic Acids in Water[J]. *Advanced Synthesis & Catalysis*, 2002. 344: 270 - 273.
- [49].Iimura S., Manabe K., and Kobayashi S.. Catalytic deprotection of protected alcohols in water using low-loading and alkylated polystyrene-supported sulfonic acid[J]. *Journal of Organic Chemistry*, 2003. 68(22): 8723-8725.

- [50].Molinari H., Montanari F., and Tundo P. Heterogeneous phase-transfer catalysts: high efficacy of catalysts bonded by a long chain to a polymer matrix[J]. *Journal of the Chemical Society, Chemical Communications*, 1977(18): 639-641.
- [51].Gruttadauria M., et al. Hydrophobically Directed Aldol Reactions: Polystyrene-Supported L-Proline as a Recyclable Catalyst for Direct Asymmetric Aldol Reactions in the Presence of Water[J]. *European Journal of Organic Chemistry*, 2007. 2007(28): 4688-4698.
- [52].Gravert D. J. and Janda K. D. Organic synthesis on soluble polymer supports: Liquid-phase methodologies[J]. *Chemical Reviews*, 1997. 97(2): 489-509.
- [53].Benaglia M, et al. Poly(ethylene glycol)-supported proline: A versatile catalyst for the enantioselective aldol and iminoaldol reactions[J]. *Advanced Synthesis & Catalysis*, 2002. 344(5): 533-542.
- [54].Castejon H. and Wiberg K. B. Solvent effects on methyl transfer reactions. 1. The Menshutkin Reaction[J]. *Journal of the American Chemical Society*, 1999. 121(10): 2139-2146.
- [55].Bond G, et al. Friedel-Crafts acylation reactions using heterogeneous catalysts stimulated by conventional and microwave heating[J]. *Journal of Molecular Catalysis a-Chemical*, 2007. 278(1-2): 1-5.
- [56].David E., Bergbreiter J T, and Chayanant H. Using Soluble Polymer Supports To Facilitate Homogeneous Catalysis[J]. *Chemical Reviews*, 2009. 109(2): 530 – 582.
- [57].Chen J., et al. A review: Non-cross-linked polystyrene bound reagents, catalysts and syntheses[J]. *Reactive and Functional Polymers*, 2006. 66(12): 1434-1451.
- [58].Li J, et al. Pd Nanoparticles in Ionic Liquid Brush: A Highly Active and Reusable Heterogeneous Catalytic Assembly for Solvent-Free or On-Water Hydrogenation of Nitroarene under Mild Conditions[J]. *ACS Catalysis*, 2011. 1(6): 657-664.
- [59].Chan M W. Weak attractive ligand-polymer and related interactions in catalysis and reactivity: impact, applications, and modeling[J]. *Chemistry-An Asian Journal*, 2008. 3(1): 18-27.
- [60].沙昆原,杨瑞华. 高分子-金属络合催化中的高分子效应[J]. *石油化工*, 1980. 5: 003.
- [61].Rana S. Influence of Resin Cross-Linking on Solid-Phase Chemistry[J]. *Journal of Combinatorial Chemistry*, 2001. 3: 9-15.
- [62].Kenneth A., Kun R K. The Pore Structure of Macroreticular Ion exchange resin[J]. *Journal of Polymer Science Part C*, 1967(16): 1457-1469.
- [63].Gruttadauria M, et al. Novel Prolinamide-Supported Polystyrene as Highly Stereoselective and Recyclable Organocatalyst for the Aldol Reaction[J]. *Advanced Synthesis & Catalysis*, 2008. 350(9): 1397-1405.
- [64].Kouzu M., Nakagaito A., and Hidaka J. S. Pre-esterification of FFA in plant oil transesterified into biodiesel with the help of solid acid catalysis of sulfonated cation-exchange resin[J]. *Applied Catalysis A: General*, 2011. 405(1-2): 36-44.
- [65].Rios L A, et al. Resin catalyzed alcoholysis of epoxidized fatty esters: Effect of the alcohol and the resin structures[J]. *Applied Catalysis A: General*, 2005. 284(1-2): 155-161.
- [66].Xu Y., Gin D. L., and Elliott B. J. Catalyzed dioctyl phthalate formation using a nanostructured solid acid resin[J]. *AIChE Journal*, 2006. 52(1): 418-421.
- [67].Wu H. S. and Lin H. J. Catalytic kinetics of quaternary ammonium poly(styrene-co- methylstyrene) resin[J]. *AIChE Journal*, 2006. 52(7): 2535-2544.
- [68].Jana S, Nath S., Pande S., et al. Synthesis of silver nanoshell-coated cationic polystyrene beads: A solid phase catalyst for the reduction of 4-nitrophenol[J]. *Applied Catalysis A: General*, 2006. 313(1): 41-48.
- [69].Price K E. Microencapsulated Linear Polymers: “ Soluble ” Heterogeneous Catalysts[J]. *Journal of the American Chemical Society*, 2006. 128: 10376-10377.
- [70].Yoshida M. Structural Control of Core-Shell Polystyrene Microcapsule-Immobilized Microbial Cells and Their Application to Polymeric Microbioreactors[J]. *Journal of Applied Polymer Science*, 2003. 89: 1966-1975.
- [71].Corma A and Garcia H. Lewis acids: From conventional homogeneous to green homogeneous and heterogeneous catalysis[J]. *Chemical Reviews*, 2003. 103(11): 4307-4366.
- [72].Mestres R. A green look at the aldol reaction[J]. *Green Chemistry*, 2004. 6(12): 583-603.

- [73].List B., and Carlos F. Barbas. Proline-Catalyzed Direct Asymmetric Aldol Reactions.[J]. Journal of the American Chemical Society, 2000. 122(10): 2395-2396.
- [74].Kolfenbach J. J.. Production of Methyl Vinyl Ketone from Methyl vinyl carbinol[J]. Industrial & Engineering Chemistry Research, 1945. 37: 1178-1180.
- [75].Qian H., Liu D. B., and Lv C. X. Synthesis of Chalcones via Claisen-Schmidt Reaction Catalyzed by Sulfonic Acid-Functional Ionic Liquids[J]. Industrial & Engineering Chemistry Research, 2011. 50(2): 1146-1149.
- [76].Feng L. C., et al. Highly efficient chemoselective construction of 2,2-dimethyl-6-substituted 4-piperidones via multi-component tandem Mannich reaction in ionic liquids[J]. Green Chemistry, 2010. 12(6): 949-952.
- [77].Zumbansen K., Dohring A., and List B. Morpholinium Trifluoroacetate-Catalyzed Aldol Condensation of Acetone with both Aromatic and Aliphatic Aldehydes[J]. Advanced Synthesis & Catalysis, 2010. 352(7): 1135-1138.
- [78].Suzuki E. and Ono Y. Aldol Condensation Reaction between Formaldehyde and Acetone over heat-treated Synthetic Hydrotalcite and Hydrotalcite-Like Compounds[J]. Bull.Chem.Soc.JPN, 1988. 61: 1008-1010.
- [79].胡微. 碱性阴离子交换树脂催化合成甲基乙烯基酮[J]. 江苏化工, 2003. 31(002): 47-48.
- [80].顾起鹤,戚蕴石. 甲醛丙酮催化缩合合成甲基乙烯基酮[J]. 石油化工, 1991. 20(1): 36-41.
- [81].徐国兴,孔识卫和陈春峰. KF/Al<sub>2</sub>O<sub>3</sub> 催化合成甲基乙烯基酮的研究[J]. 山东化工, 2006. 35(5): 1-3.
- [82].Okada M. and Asami Y. Selective Formation of Methyl Vinyl Ketone From Acetone and Methanol With Silical-Magnesia Base Catalyst[J]. Chemistry Letters, 1973. 2(4): 333-334.
- [83].陈德安,欧光南,翁玉攀. 由甲醇和丙酮合成甲基乙烯基酮铜基催化剂的研究[J]. 厦门大学学报, 1994. 33(4): 481-487.
- [84].Tanne R and P Gill. Aldol condensation reactions of acetone and formaldehyde over vanadium phosphate catalysts: Comments on the acid – base properties[J]. Phys. Chem. Chem. Phys., 2002. 4: 688-695.
- [85].Fajardo H V. Gas-Phase Selective Conjugate Addition of Methanol to Acetone for Methyl Vinyl Ketone over SnO<sub>2</sub> Nanoparticle Catalysts[J]. J. Braz. Chem. Soc., 2005. 16: 607-613.
- [86].Gelbard G. Organic Synthesis by Catalysis with Ion-Exchange Resins[J]. Industrial & Engineering Chemistry Research, 2005. 44: 8468-8498.
- [87].Zlotin S. G., Kucherenko A. S., and Beletskaya I. P. Organocatalysis of asymmetric aldol reaction. Catalysts and reagents[J]. Russian Chemical Reviews, 2009. 78(8): 737 - 784.
- [88].Luo S, et al. Noncovalently supported heterogeneous chiral amine catalysts for asymmetric direct aldol and Michael addition reactions[J]. Chemistry, 2008. 14(4): 1273-1281.
- [89].Susumu Saito H Y. Design of Acid-Base Catalysis for the Asymmetric Direct Aldol Reaction[J]. Accounts of Chemical Research, 2004. 37: 570-579.
- [90].Serra-Holm T S. Aldolization of butyraldehyde with formaldehyde over a commercial anion exchange resin kinetics and selectivity aspects[J]. Applied Catalysis A: General, 2000. 198: 207-221.
- [91].Serra-Holm T S. Comparison of Activity and Selectivity of Weakly Basic Anion-Exchange catalysts for the aldolization of butyraldehyde with formaldehyde[J]. Organic Process Research & Development, 2001. 5: 368-375.
- [92].Susumu S. Design of Acid-Base Catalysis for the Asymmetric Direct Aldol Reaction[J]. Accounts of Chemical Research, 2004. 37: 570-579.
- [93].Barbaro P. and Liguori F. Ion Exchange Resins: Catalyst Recovery and Recycle[J]. Chemical Reviews, 2009. 109(2): 515-529.
- [94].Cozzi F. Immobilization of Organic Catalysts: When, Why, and How[J]. Advanced Synthesis & Catalysis, 2006. 348(12-13): 1367-1390.
- [95].Podrebarac. A kinetic study of the aldol condensation of acetone using an anion exchange resin catalyst[J]. Chemical Engineering Science, 1997. 52(17): 2991-3002.
- [96].Astle J. M. and Zaslowaky J. A. Reactions Catalyzed by Anion Exchange Resins--aldol condensation[J]. Industrial and Engineering Chemistry, 1952. 44(12): 2869-2871.

- [97]. Rao Q., Luo S., and Gong L. Water as dual functional cocatalyst: A theoretical study on the mechanism of direct aldol reaction on water catalyzed by a leucine derivative[J]. Chinese Science Bulletin, 2010. 55(17): 1742-1752.
- [98]. Girolamo M and Marchionna M. Acidic and basic ion exchange resins for industrial applications[J]. Journal of Molecular Catalysis A: Chemical, 2001. 177(1): 33-40.
- [99]. Sanchez C., et al. Applications of advanced hybrid organic-inorganic nanomaterials: from laboratory to market[J]. Chemical Society Reviews, 2011. 40(2): 696-753.
- [100]. 何仁, 陶晓春, 张兆国. 金属有机化学[M]. 化学与应用化学研究生教学用书. 上海: 华东理工大学出版社. 2007.
- [101]. McNamara C. A., Dixon M. J., and Bradley M. Recoverable Catalysts and Reagents Using Recyclable Polystyrene-Based Supports[J]. Chemical Reviews, 2002. 102(10): 3275-3300.
- [102]. Jang S-B. Polymer-bound palladium-catalyzed cross-coupling of organoboron compounds with organic halides and organic triflates[J]. Tetrahedron letters, 1997. 38(10): 1793-1796.
- [103]. Fenger I. and Drian C. Le. Reusable polymer-supported palladium catalysts: An alternative to tetrakis (triphenylphosphine) palladium in the Suzuki cross-coupling reaction[J]. Tetrahedron letters, 1998. 39(24): 4287-4290.
- [104]. Dygutsch D. P. and Eilbracht P. Synthesis of cyclopentanone derivatives with polystyrene -supported cyclopentadienyl rhodium catalysts[J]. Tetrahedron, 1996. 52(15): 5461-5468.
- [105]. Lee J. H., Lee Y., and Nam J. D. Multi-metal shell coated with surface modification of polystyrene bead cores to improve its adhesion metal shell[J]. Surface and Interface Analysis, 2009: 36 – 39.
- [106]. Yang Y, Lee Y., Nam J. D. et al. Polystyrene-ZnO core-shell microspheres and hollow ZnO structures synthesized with the sulfonated polystyrene templates[J]. Journal of Solid State Chemistry, 2006. 179(2): 470-475.
- [107]. Schroder M. Functional Metal-Organic Frameworks: Gas Storage, Separation and Catalysis[M]. Springer-Verlag Berlin Heidelberg. 75-206. 2010.
- [108]. Farrusseng D. Metal-Organic Frameworks: Applications from Catalysis to Gas Storage [M]. Wiley-VCH Verlag GMBH & Co. KGaA. 191-210. 2011.
- [109]. Macgillivray L R. Metal-Organic Frameworks: Design and Application[M]. John Wiley & Sons. Inc. 37-90. 2010.
- [110]. 魏文英. 新型金属有机骨架的合成, 结构表征及催化性能[D]. 天津大学. 2005.
- [111]. Kleist W., et al. Mixed-Linker Metal-Organic Frameworks as Catalysts for the Synthesis of Propylene Carbonate from Propylene Oxide and CO<sub>2</sub>[J]. European Journal of Inorganic Chemistry, 2009. 2009(24): 3552-3561.
- [112]. Paz F A, et al. Ligand design for functional metal-organic frameworks[J]. Chemical Society Reviews, 2012. 41(3): 1088-1110.
- [113]. Fang Q R, et al. Functional mesoporous metal-organic frameworks for the capture of heavy metal ions and size-selective catalysis[J]. Inorg Chem, 2010. 49(24): 11637-11642.
- [114]. Xuan W, et al. Mesoporous metal-organic framework materials[J]. Chemical Society Reviews, 2012. 41(5): 1677-1695.
- [115]. Zacher D, et al. Surface chemistry of metal-organic frameworks at the liquid-solid interface[J]. Angewandte Chemie International Edition, 2011. 50(1): 176-199.
- [116]. Füllgel E A, et al. Synthetic routes toward MOF nanomorphologies[J]. Journal of Materials Chemistry, 2012. 22(20): 10119-10133.
- [117]. Cohen S M. Postsynthetic methods for the functionalization of metal-organic frameworks[J]. Chem Rev, 2012. 112(2): 970-1000.
- [118]. Schröder F. and Fischer R. A. Doping of Metal-Organic Frameworks with Functional Guest Molecules and Nanoparticles[J]. Topic Current Chemistry, 2010. 293: 77-113.
- [119]. Yuan B, et al. A Highly Active Heterogeneous Palladium Catalyst for the Suzuki – Miyaura and Ullmann

- Coupling Reactions of Aryl Chlorides in Aqueous Media[J]. *Angewandte Chemie International Edition*, 2010. 49(24): 4054-4058.
- [120].Hwang Y K, et al. Amine grafting on coordinatively unsaturated metal centers of MOFs: consequences for catalysis and metal encapsulation[J]. *Angewandte Chemie International Edition*, 2008. 47(22): 4144-4148.
- [121].Lu G, Guo Z., Farha O. K. et al. Imparting functionality to a metal-organic framework material by controlled nanoparticle encapsulation[J]. *Nature Chemistry*, 2012. 4(4): 310-316.
- [122].Zhan W. W., et al. Semiconductor@Metal – Organic Framework Core – Shell Heterostructures: A Case of ZnO@ZIF-8 Nanorods with Selective Photoelectrochemical Response[J]. *Journal of the American Chemical Society*, 2013. 135(5): 1926-1933.
- [123].Jiang H. L., Ishida T., Haruta M., et al. Synergistic catalysis of Au@Ag core-shell nanoparticles stabilized on metal-organic framework[J]. *Journal of the American Chemical Society*, 2011. 133(5): 1304-1306.
- [124].Liu H. Y., et al. Solvothermal Assembly of a Series of Organic – Inorganic Hybrid Materials Constructed from Keggin Polyoxometalate Clusters and Copper(I) – Organic Frameworks[J]. *Crystal Growth & Design*, 2011. 11(5): 1786-1797.
- [125].Müller M., et al. Highly emissive metal – organic framework composites by host – guest chemistry[J]. *Photochemical & Photobiological Sciences*, 2010. 9(6): 846-853.
- [126].Wu C. D., et al. A homochiral porous metal-organic framework for highly enantioselective heterogeneous asymmetric catalysis[J]. *Journal of the American Chemical Society*, 2005. 127(25): 8940-8941.
- [127].Wang Z. and Cohen S. M. Postsynthetic covalent modification of a neutral metal-organic framework[J]. *Journal of the American Chemical Society*, 2007. 129(41): 12368-12369.
- [128].Xiang Z., Cao D., Yang W.T., et al. Metal-organic frameworks with incorporated carbon nanotubes: improving carbon dioxide and methane storage capacities by lithium doping[J]. *Angewandte Chemie International Edition*, 2011. 50(2): 491-494.
- [129].Petit C. and Bandoz T. J. MOF-Graphite Oxide Composites: Combining the Uniqueness of Graphene Layers and Metal-Organic Frameworks[J]. *Advanced Materials*, 2009. 21: 4753 – 4757.
- [130].Yu L. Q. and Yan X. P. Covalent bonding of zeolitic imidazolate framework-90 to functionalized silica fibers for solid-phase microextraction[J]. *Chemical Communications*, 2013. 49(21): 2142-2144.
- [131].Karimi Z. and Morsali A. Modulated formation of metal-organic frameworks by oriented growth over mesoporous silica[J]. *Journal of Materials Chemistry A*, 2013. 1(9): 3047-3054.
- [132].Jin Y, et al. Microwave-assisted syntheses of highly CO<sub>2</sub>-selective organic cage frameworks (OCFs)[J]. *Chemical Science*, 2012. 3(3): 874-877.
- [133].Lan J., Wang W.C., Smit B., et al. Doping of Alkali, Alkaline-Earth, and Transition Metals in Covalent-Organic Frameworks for Enhancing CO<sub>2</sub> Capture by First-Principles Calculations and Molecular Simulations[J]. *ACS Nano*, 2010. 4(7): 4225 – 4237.
- [134].Farrusseng D., Aguado S., and Pinel C. Metal-organic frameworks: opportunities for catalysis[J]. *Angewandte Chemie International Edition*, 2009. 48(41): 7502-7513.
- [135].Li J. R., Sculley J., and Zhou H. C. Metal-organic frameworks for separations[J]. *Chemical Reviews*, 2012. 112(2): 869-932.
- [136].Furukawa H. and Yaghi O. M. Storage of Hydrogen, Methane, and Carbon Dioxide in Highly Porous Covalent Organic Frameworks for Clean Energy Applications[J]. *Journal of the American Chemical Society*, 2009. 131: 8875 – 8883.
- [137].Betard A. and Fischer R. A. Metal-organic framework thin films: from fundamentals to applications[J]. *Chemical Reviews*, 2012. 112(2): 1055-1083.
- [138].Jiang H. L. and Xu Q. Porous metal-organic frameworks as platforms for functional applications[J]. *Chemical Communications*, 2011. 47(12): 3351-3370.
- [139].尹作娟等. 金属有机骨架的合成及应用[J]. *化学与粘合*, 2009(003): 61-65.
- [140].Cui Y. J., Qian G. D. and Chen B. L. Luminescent functional metal-organic frameworks[J]. *Chemical Reviews*, 2012. 112(2): 1126-1162.

- [141].Zhang W. and Xiong R. G. Ferroelectric metal-organic frameworks[J]. Chemical Reviews, 2012. 112(2): 1163-1195.
- [142].Kreno L. E., et al. Metal-organic framework materials as chemical sensors[J]. Chemical Reviews, 2012. 112(2): 1105-1125.
- [143].Horcajada P, et al. Metal-organic frameworks in biomedicine[J]. Chemical Reviews, 2012. 112(2): 1232-1268.
- [144].Ma Z. and Moulton B. Recent advances of discrete coordination complexes and coordination polymers in drug delivery[J]. Coordination Chemistry Reviews, 2011. 255(15-16): 1623-1641.
- [145].Czaja A. U., Trukhan N, and Muller U. Industrial applications of metal-organic frameworks[J]. Chemical Society Reviews, 2009. 38(5): 1284-1293.
- [146].Kondo M, et al. Microporous materials constructed from the interpenetrated coordination networks. Structures and methane adsorption properties[J]. Chemistry of materials, 2000. 12(5): 1288-1299.
- [147].Yang J., Wolvertonb C. and Siegel D. J. High capacity hydrogen storage materials: attributes for automotive applications and techniques for materials discovery[J]. Chemical Society Reviews, 2010. 39(2): 656-675.
- [148].Suh M. P., Prasad T.K. and Lim D.W. Hydrogen storage in metal-organic frameworks[J]. Chemical Reviews, 2012. 112(2): 782-835.
- [149].Getman R. B., Wilmer C. E. and Snurr R. Q. Review and analysis of molecular simulations of methane, hydrogen, and acetylene storage in metal-organic frameworks[J]. Chemical Reviews, 2012. 112(2): 703-723.
- [150].Dybtsev D. N., et al. Microporous manganese formate: a simple metal-organic porous material with high framework stability and highly selective gas sorption properties[J]. Journal of the American Chemical Society, 2004. 126(1): 32-33.
- [151].Meek S. T., Greathouse J. A., and Allendorf M. D. Metal-organic frameworks: a rapidly growing class of versatile nanoporous materials[J]. Advanced Materials, 2011. 23(2): 249-267.
- [152].Wu H., Olson D. H., and Li J. Commensurate adsorption of hydrocarbons and alcohols in microporous metal organic frameworks[J]. Chemical Reviews, 2012. 112(2): 836-868.
- [153].Li J. R., et al. Carbon dioxide capture-related gas adsorption and separation in metal-organic frameworks[J]. Coordination Chemistry Reviews, 2011. 255(15-16): 1791-1823.
- [154].Liu J., McGrail B. P., Brown D. R., et al. Progress in adsorption-based CO<sub>2</sub> capture by metal-organic frameworks[J]. Chemical Society Reviews, 2012. 41(6): 2308-2322.
- [155].Keskin S., Heest T. M. V., and Sholl D. S. Can metal-organic framework materials play a useful role in large-scale carbon dioxide separations?[J]. ChemSusChem, 2010. 3(8): 879-891.
- [156].Duan J. G., Bai J. F., Zheng B. S., et al. Highly selective CO<sub>2</sub> capture of an agw-type metal-organic framework with inserted amides: experimental and theoretical studies[J]. Chemical Communications, 2012. 48(25): 3058-3060.
- [157].Sumida K, Mason J. A., McDonald T. M., et al. Carbon dioxide capture in metal-organic frameworks[J]. Chemical Reviews, 2012. 112(2): 724-781.
- [158].D'Alessandro D. M., Smit B., and Long J. R.. Carbon dioxide capture: prospects for new materials[J]. Angewandte Chemie International Edition. 49(35): 6058-6082.
- [159].Kuppler R J, Fang Q. R., Li J. R. et al. Potential applications of metal-organic frameworks[J]. Coordination Chemistry Reviews, 2009. 253: 3042 – 3066.
- [160].Jones C. W. and Maginn E. J. Materials and processes for carbon capture and sequestration[J]. ChemSusChem, 2010. 3(8): 863-864.
- [161].Keskin S., Heest T. M. and Sholl D. S. Can Metal – Organic Framework Materials Play a Useful Role in Large Scale Carbon Dioxide Separations?[J]. ChemSusChem, 2010. 3(8): 879-891.
- [162].Bae Y. S. and Snurr R. Q. Development and evaluation of porous materials for carbon dioxide separation and capture[J]. Angewandte Chemie International Edition, 2011. 50(49): 11586-11596.
- [163].Wang B., Furukawa H., Keffe M., et al. Colossal cages in zeolitic imidazolate frameworks as selective



carbon dioxide reservoirs[J]. *Nature*, 2008. 453(7192): 207-211.

[164].Rahul B., David B., Carolyn K., et al. Control of Pore Size and Functionality in Isostructural Zeolitic Imidazolate Frameworks and their Carbon Dioxide Selective Capture Properties[J]. *Journal of the American Chemical Society*, 2009. 131: 3875 – 3877.

[165].Phan A., Romo F. J., Knobler C. B., et al. Synthesis, Structure, and Carbon Dioxide Capture Properties of Zeolitic Imidazolate Frameworks[J]. *Accounts of Chemical Research*, 2010. 43(1): 58-67.

[166].Sabouni R., Kazemian H., and Rohani S. Carbon dioxide adsorption in microwave-synthesized metal organic framework CPM-5: Equilibrium and kinetics study[J]. *Microporous and Mesoporous Materials*, 2013. 175: 85-91.

[167].Yao Q., et al. Interpenetrated metal – organic frameworks and their uptake of CO<sub>2</sub> at relatively low pressures[J]. *Journal of Materials Chemistry*, 2012. 22(20): 10345 – 10351.

[168].Choi Y. J., et al. Covalent organic frameworks for extremely high reversible CO<sub>2</sub> uptake capacity: a theoretical approach[J]. *Journal of Materials Chemistry*, 2011. 21(4): 1073 – 1078.

[169].Wu H. and Yildirim T. Adsorption Sites and Binding Nature of CO<sub>2</sub> in Prototypical Metal-Organic Frameworks: A Combined Neutron Diffraction and First-Principles Study[J]. *The Journal of Physical Chemistry Letters*, 2010. 1: 1946 – 1951.

[170].Chen Y. F., Eddaoudi M., and Jiang J. W. CO<sub>2</sub> adsorption in mono-, di- and trivalent cation-exchanged metal-organic frameworks: a molecular simulation study[J]. *Langmuir*, 2012. 28(8): 3903-3910.

[171].Park J., et al. Tuning Metal – Organic Frameworks with Open-Metal Sites and Its Origin for Enhancing CO<sub>2</sub> Affinity by Metal Substitution[J]. *The Journal of Physical Chemistry Letters*, 2012. 3(7): 826-829.

[172].Gupta K. M., Hu Z. Q. and Jiang J. W. Metal-organic framework supported ionic liquid membranes for CO<sub>2</sub> capture: anion effects[J]. *Physical Chemistry Chemical Physics*, 2012. 14(16): 5785-5794.

[173].Vogiatzis K. D., Kloppe W. and Froudakis. G. E. Ab initio study of the interactions between CO<sub>2</sub> and N-containing organic heterocycles[J]. *ChemPhysChem*, 2009. 10(2): 374-383.

[174].An J., Geib S. J. and Rosi N. L. Synthesis, Structure, Assembly, and Modulation of the CO<sub>2</sub> Adsorption Properties of a Zinc-Adeninate Macrocyclic[J]. *Journal of the American Chemical Society*, 2009. 131: 8401 – 8403.

[175].An J., Geib S. J., and Rosi N. L. High and Selective CO<sub>2</sub> Uptake in a Cobalt Adeninate Metal-Organic Framework Exhibiting Pyrimidine- and Amino-Decorated Pores[J]. *Journal of the American Chemical Society*, 2010. 132: 38 – 39.

[176].Wang F., Yang H., Zhang H. X., et al. A new approach towards tetrahedral imidazolate frameworks for high and selective CO<sub>2</sub> uptake[J]. *Chemical Communications*, 2011. 47(20): 5828-5830.

[177].An J. Porous Metal-Adeninate Crystalline Materials: Design, Synthesis and Emerging Properties[D]. Doctor of Philosophy. University of Pittsburgh. 242. 2011.

[178].Qin J. S., et al. N-rich zeolite-like metal – organic framework with sodalite topology: high CO<sub>2</sub> uptake, selective gas adsorption and efficient drug delivery[J]. *Chemical Science*, 2012. 3(6): 2114-2118.

[179].Zhang J. P., Lin J. B., and Chen X. M. Metal azolate frameworks: from crystal engineering to functional materials[J]. *Chemical Reviews*, 2012. 112(2): 1001-1033.

[180].William M., Furukawa H., Yaghi O. K., et al. A Combined Experimental&#8722;Computational Investigation of Carbon Dioxide[J]. *Journal of the American Chemical Society*, 2010. 132: 11006 – 11008.

[181].Demessence A., Foo M. L., and Long J. R. Strong CO<sub>2</sub> Binding in a Water-Stable, Triazolate-Bridged Metal&#8722;Organic Framework Functionalized with Ethylenediamine[J]. *Journal of the American Chemical Society*, 2009. 131: 8784 – 8786.

[182].Vaidhyanathan R., Dawson K. W. and Shimizu G. An amine-functionalized metal organic framework for preferential CO<sub>2</sub> adsorption at low pressures[J]. *Chemical Communications*, 2009(35): 5230-5232.

[183].Das A., et al. Application of the piperazine-grafted CuBTTri metal-organic framework in postcombustion carbon dioxide capture[J]. *Microporous and Mesoporous Materials*, 2013. 174: 74-80.

[184].Planas N, Poloni R., Lin L. C., et al. The mechanism of carbon dioxide adsorption in an alkylamine-

functionalized metal-organic framework[J]. *Journal of the American Chemical Society*, 2013. 135(20): 7402-7405.

[185]. Babarao R., Dai S., and Jiang D. E. Functionalizing porous aromatic frameworks with polar organic groups for high-capacity and selective CO<sub>2</sub> separation: a molecular simulation study[J]. *Langmuir*, 2011. 27(7): 3451-3460.

[186]. Liu X, et al. Enhanced carbon dioxide uptake by metalloporphyrin-based microporous covalent triazine framework[J]. *Polymer Chemistry*, 2013. 4(8): 2445 – 2448.

[187]. Mueller U, et al. Organometallic building materials and method for producing the same. 2006, Patents.

[188]. Snejko N, Gomez L. B., Guti  rrez P., et al. From rational octahedron design to reticulation serendipity. A thermally stable rare earth polymeric disulfonate family with CdI<sub>2</sub>-like structure, bifunctional catalysis and optical properties[J]. *Chemical Reviews*, 2002(13): 1366-1367.

[189]. Ranocchi M and Bokhoven J. A. Catalysis by metal – organic frameworks: fundamentals and opportunities[J]. *Physical Chemistry Chemical Physics*, 2011. 13(14): 6373 – 6712.

[190]. Chui S. S. Y, et al. A chemically functionalizable nanoporous material [Cu<sub>3</sub>(TMA)<sub>2</sub>(H<sub>2</sub>O)<sub>3</sub>]<sub>n</sub> [J]. *Science*, 1999. 283(5405): 1148-1150.

[191]. Gascon J, Hernandez-Alonso M. D., Gerard P.M., et al. Amino-based metal-organic frameworks as stable, highly active basic catalysts[J]. *Journal of Catalysis*, 2009. 261(1): 75-87.

[192]. Dhakshinamoorthy A. and Garcia H. Catalysis by metal nanoparticles embedded on metal-organic frameworks[J]. *Chemical Society Reviews*, 2012. 41(15): 5262-5284.

[193]. Uemura T., et al. Nanochannel Promoted Polymerization of Substituted Acetylenes in Porous Coordination Polymers[J]. *Angewandte Chemie International Edition*, 2006. 118(25): 4218-4222.

[194]. Lee J., et al. Metal-organic framework materials as catalysts[J]. *Chemical Society Reviews*, 2009. 38(5): 1450-1459.

[195]. Corma A, Garc  a H., and Xamena F. X. Engineering Metal Organic Frameworks for Heterogeneous Catalysis.[J]. *Chemical Reviews*, 2010. 110(8): 4606 – 4655.

[196]. Aresta M. Carbon Dioxide as Chemical Feedstock[M]. Weinheim: WILEY-VCH Verlag GmbH & Co. KGaA. 2010.

[197]. Quadrelli E. A., Duplan J. L. and Perathoner S. Carbon dioxide recycling: emerging large-scale technologies with industrial potential[J]. *ChemSusChem*, 2011. 4(9): 1194-1215.

[198]. Chu S. Carbon Capture and Sequestration[J]. *Science*, 2009. 325: 1599-1621.

[199]. 韩玉杰. CO<sub>2</sub>的捕集和封存技术的现状, 前景与挑战[J]. *应用能源技术*, 2009(006): 38-40.

[200]. 王献红. 二氧化碳的固定与利用[M]. 北京 化学工业出版社. 2011.

[201]. Jutz F., and Baiker A.. Ionic Liquids and Dense Carbon Dioxide: A Beneficial Biphasic System for Catalysis[J]. *Chemical Reviews*, 2011. 111(2): 322 – 353.

[202]. Solinas D. G. Beneficial Combination of Ionic Liquids and Carbon Dioxide in Transition Metal Catalysed Reactions.[J]. *Current Organic Chemistry*, 2009. 13: 1300-1321.

[203]. Trost B. M. The atom economy-a search for synthetic efficiency[J]. *Science*, 1991. 254(5037): 1471-1477.

[204]. Peters M., Kuckshinrichs W., Leitner W., et al. Chemical technologies for exploiting and recycling carbon dioxide into the value chain[J]. *ChemSusChem*, 2011. 4(9): 1216-1240.

[205]. Huang K., Sunw C. L., and Shi Z. J. Transition-metal-catalyzed C-C bond formation through the fixation of carbon dioxide[J]. *Chemical Society Reviews*, 2011. 40(5): 2435-2452.

[206]. Sakakura T., Choi J. C, and Yasuda H. Transformation of Carbon Dioxide[J]. *Chemical Reviews*, 2007. 107(6): 2365-2387.

[207]. Gibson D. H. The organometallic chemistry of carbon dioxide[J]. *Chemical Reviews*, 1996. 96(6): 2063-2096.

[208]. Leitner W. The coordination chemistry of carbon dioxide and its relevance for catalysis: a critical survey[J]. *Coordination Chemistry Reviews*, 1996. 153: 257-284.

[209]. Yin X. and Moss J. R. Recent developments in the activation of carbon dioxide by metal complexes[J].

Coordination Chemistry Reviews, 1999. 181(1): 27-59.

[210].Braunstein P., Matt D., and Nobel D. Reactions of Carbon Dioxide with Carbon-Carbon Bond Formation Catalyzed by Transition-Metal Complexes[J]. Chemical Reviews, 1988. 88(5): 747-764.

[211].Huff C. A. and Sanford M. S. Cascade catalysis for the homogeneous hydrogenation of CO<sub>2</sub> to methanol[J]. Journal of the American Chemical Society, 2011. 133(45): 18122-18125.

[212].Dixneuf P. H. Bifunctional catalysis: a bridge from CO<sub>2</sub> to methanol[J]. Nature Chemistry, 2011. 3(8): 578-579.

[213].Sakakura T. and Kohno K. The synthesis of organic carbonates from carbon dioxide[J]. Chemical Communications, 2009(11): 1312-1330.

[214].Clements J. H. Reactive applications of cyclic alkylene carbonates[J]. Industrial & Engineering Chemistry Research, 2003. 42(4): 663-674.

[215].Shaikh A. A. G. and Sivaram S. Organic carbonates[J]. Chemical Reviews, 1996. 96(3): 951-976.

[216].Sun J., Fujita S., and Arai M. Development in the green synthesis of cyclic carbonate from carbon dioxide using ionic liquids[J]. Journal of organometallic chemistry, 2005. 690(15): 3490-3497.

[217].Sakakura T. and Kohno K. The synthesis of organic carbonates from carbon dioxide[J]. Chemical Communications, 2009(11): 1312-1330.

[218].Eta V., et al. Synthesis of Dimethyl Carbonate from Methanol and Carbon Dioxide: The Effect of Dehydration. Available from: <http://archivos.labcontrol.cl/wcce8/offline/techsched/manuscripts/6dmz95.pdf>.

[219].Honda M, Noorjahan B., Fujimoto K., et al. Low pressure CO<sub>2</sub> to dimethyl carbonate by the reaction with methanol promoted by acetonitrile hydration[J]. Chemical Communications. 2009(30): 4596-4598.

[220].Choi J. C., et al. Tin- or titanium-catalyzed dimethyl carbonate synthesis from carbon dioxide and methanol: Large promotion by a small amount of triflate salts[J]. Catalysis Communications, 2008. 9(7): 1630-1633.

[221].Kohno K., Ohshima Y., Yasuda H., et al. Synthesis of dimethyl carbonate from carbon dioxide catalyzed by titanium alkoxides with polyether-type ligands[J]. ChemSusChem, 2008. 1(3): 186-188.

[222].Choi J. C., et al. Selective and high yield synthesis of dimethyl carbonate directly from carbon dioxide and methanol[J]. Green Chemistry, 2002. 4(3): 230-234.

[223].Zhang Z. F., et al. Synthesis of Dimethyl Carbonate from Carbon Dioxide and Methanol over CexZr1-xO<sub>2</sub> and [EMIM]Br/Ce0.5Zr0.5O<sub>2</sub>[J]. Industrial & Engineering Chemistry Research, 2011. 50(4): 1981-1988.

[224].Cai Q., Guo L. J. and Shan Y. K.. Studies on synthesis of dimethyl carbonate from methanol and carbon dioxide[J]. Catalysis Communications, 2009. 10(5): 605-609.

[225].Fang S. N. Direct synthesis of dimethyl carbonate from carbon dioxide and methanol catalyzed by base[J]. Applied Catalysis A: General, 1996. 142: 1-3.

[226].Fujita S. I., et al. Synthesis of dimethyl carbonate from carbon dioxide and methanol in the presence of methyl iodide and base catalysts under mild conditions: effect of reaction conditions and reaction mechanism[J]. Green Chemistry, 2001. 3(2): 87-91.

[227].Eta V., Leino A. R., Kordas K., et al. Synthesis of Dimethyl Carbonate from Methanol and Carbon Dioxide: Circumventing Thermodynamic Limitations[J]. Industrial & Engineering Chemistry Research. 2010. 49: 9609 – 9617.

[228].Hou Z., et al. Synthesis of dimethyl carbonate using CO<sub>2</sub> and methanol: enhancing the conversion by controlling the phase behavior[J]. Green Chemistry, 2002. 4(5): 467-471.

[229].Lee M. H., et al. Measurement of VLE data of carbon dioxide+dimethyl carbonate system for the direct synthesis of dimethyl carbonate using supercritical CO<sub>2</sub> and methanol[J]. Fluid Phase Equilibria, 2012. 318: 77-82.

[230].Zhou Y., et al. Novel Cu – Fe bimetal catalyst for the formation of dimethyl carbonate from carbon dioxide and methanol[J]. RSC Advances, 2012. 2(17): 6831-6838.

[231].Tomishige K., Ikeda Y. and Fujimoto K.. A novel method of direct synthesis of dimethyl carbonate from methanol and carbon dioxide catalyzed by zirconia[J]. Catalysis Letters, 1999. 58: 225 – 229.

- [232].Aresta M., et al. Comparison of the behaviour of supported homogeneous catalysts in the synthesis of dimethylcarbonate from methanol and carbon dioxide: Polystyrene-grafted tin-metallorganic species versus silesquioxanes linked Nb-methoxo species[J]. *Inorganica Chimica Acta*, 2008. 361(11): 3215-3220.
- [233].Kalhor M. P., et al. From CO<sub>2</sub> to dimethyl carbonate with dialkyldimethoxystannanes: the key role of monomeric species[J]. *Physical Chemistry Chemical Physics*, 2011. 13(6): 2401-2408.
- [234].Fan B., et al. Organotin compounds immobilized on mesoporous silicas as heterogeneous catalysts for direct synthesis of dimethyl carbonate from methanol and carbon dioxide[J]. *Applied Catalysis A: General*, 2010. 372(1): 94-102.
- [235].Dai W. L., et al. The direct transformation of carbon dioxide to organic carbonates over heterogeneous catalysts[J]. *Applied Catalysis A: General*, 2009. 366(1): 2-12.
- [236].Bian J., et al. Direct synthesis of dimethyl carbonate over activated carbon supported Cu-based catalysts[J]. *Chemical Engineering Journal*, 2010. 165: 686 – 692.
- [237].Tomishige K., Ikeda Y., Asadullah M. and Fujimoto K. CeO<sub>2</sub> – ZrO<sub>2</sub> solid solution catalyst for selective synthesis of dimethyl carbonate from methanol and carbon dioxide[J]. *Catalysis Letters*, 2001. 76(1-2): 71-74.
- [238].Bian J., et al. Highly effective synthesis of dimethyl carbonate from methanol and carbon dioxide using a novel copper – nickel/graphite bimetallic nanocomposite catalyst[J]. *Chemical Engineering Journal*, 2009. 147(2-3): 287-296.
- [239].La K. W., et al. Effect of acid – base properties of H<sub>3</sub>PW<sub>12</sub>O<sub>40</sub>/CeTi<sub>1-x</sub>Al<sub>x</sub>O<sub>2</sub> catalysts on the direct synthesis of dimethyl carbonate from methanol and carbon dioxide: A TPD study of H<sub>3</sub>PW<sub>12</sub>O<sub>40</sub>/CeTi<sub>1-x</sub>Al<sub>x</sub>O<sub>2</sub> catalysts[J]. *Journal of Molecular Catalysis A: Chemical*, 2007. 269(1-2): 41-45.
- [240].Almusaiteer K. Synthesis of dimethyl carbonate (DMC) from methanol and CO<sub>2</sub> over Rh-supported catalysts[J]. *Catalysis Communications*, 2009. 10(7): 1127-1131.
- [241].Bian J., et al. Carbon nanotubes supported Cu – Ni bimetallic catalysts and their properties for the direct synthesis of dimethyl carbonate from methanol and carbon dioxide[J]. *Applied Surface Science*, 2009. 255(16): 7188-7196.
- [242].Pescarmona P. P. and Taherimehr M. Challenges in the catalytic synthesis of cyclic and polymeric carbonates from epoxides and CO<sub>2</sub>[J]. *Catalysis Science & Technology*, 2012. 2(11): 2169 – 2187.
- [243].Bhanage B. M., et al. Synthesis of dimethyl carbonate and glycols from carbon dioxide, epoxides, and methanol using heterogeneous basic metal oxide catalysts with high activity and selectivity[J]. *Applied Catalysis A: General*, 2001. 219(1): 259-266.
- [244].Dokocil E. J. Effect of water and alkali modifications on ETS-10 for the cycloaddition of CO<sub>2</sub> to propylene oxide[J]. *The Journal of Physical Chemistry B*, 2005. 109(6): 2315-2320.
- [245].Kawanami H. and Ikushima Y. Chemical fixation of carbon dioxide to styrene carbonate under supercritical conditions with DMF in the absence of any additional catalysts[J]. *Chemical Communications*, 2000(21): 2089-2090.
- [246].Lu X. B., et al. Asymmetric catalysis with CO<sub>2</sub>: Direct synthesis of optically active propylene carbonate from racemic epoxides[J]. *Journal of the American Chemical Society*, 2004. 126(12): 3732-3733.
- [247].Kim H. S., et al. Imidazolium zinc tetrahalide-catalyzed coupling reaction of CO<sub>2</sub> and ethylene oxide or propylene oxide[J]. *Journal of Catalysis*, 2003. 220(1): 44-46.
- [248].Alvaro M., et al. CO<sub>2</sub> fixation using recoverable chromium salen catalysts: use of ionic liquids as cosolvent or high-surface-area silicates as supports[J]. *Journal of Catalysis*, 2004. 228(1): 254-258.
- [249].Alvaro M., et al. Polymer-bound aluminium salen complex as reusable catalysts for CO<sub>2</sub> insertion into epoxides[J]. *Tetrahedron*, 2005. 61(51): 12131-12139.
- [250].Darensbourg D. J. Making Plastics from Carbon Dioxide: Salen Metal Complexes as Catalysts for the Production of Polycarbonates from Epoxides and CO<sub>2</sub>[J]. *Chemical Reviews*, 2007. 107(6): 2388-2410.
- [251].Donald J., Darensbourg J. C. Y., Cesar O., and Fang C.. Comparative Kinetic Studies of the Copolymerization of Cyclohexene Oxide and Propylene Oxide with Carbon Dioxide in the Presence of

- Chromium Salen Derivatives. In Situ FTIR Measurements of Copolymer vs Cyclic Carbonate Production[J]. Journal of the American Chemical Society, 2003. 125: 7586-7591.
- [252]. Carly E. et al. Cobalttoporphyrin-Catalyzed CO<sub>2</sub>/Epoxide Copolymerization: Selectivity Control by Molecular Design[J]. Macromolecules, 2012. 45: 6840-6849.
- [253]. Nakano K, et al. Copolymerization of epoxides with carbon dioxide catalyzed by iron-corrole complexes: synthesis of a crystalline copolymer[J]. Journal of the American Chemical Society, 2013. 135(23): 8456-8459.
- [254]. Lindsey A. S. and Jeskey H. The Kolbe-schmitt reaction[J]. Chemical Reviews, 1957. 57(4): 583-620.
- [255]. Markovic Z., et al. Mechanism of the Kolbe-Schmitt reaction. Structure of the intermediate potassium phenoxide-CO<sub>2</sub> complex[J]. Journal of chemical information and modeling, 2007. 47(4): 1520-1525.
- [256]. Markovic Z. and Markovic S. Last step of the para route of the Kolbe-Schmitt reaction[J]. Journal of Chemical Information and Modeling, 2008. 48(1): 143-147.
- [257]. Olah G. A., et al. Efficient Chemoselective Carboxylation of Aromatics to Arylcarboxylic Acids with a Superelectrophilically Activated Carbon Dioxide-Al<sub>2</sub>Cl<sub>6</sub>/Al System [J]. Journal of the American Chemical Society, 2002. 124: 11379-11391.
- [258]. Nemoto K., et al. Direct Carboxylation of Arenes and Halobenzenes with CO<sub>2</sub> by the Combined Use of AlBr<sub>3</sub> and R<sub>3</sub>SiCl[J]. The Journal of Organic Chemistry, 2010. 75: 7855 – 7862.
- [259]. Schafer A., et al. Silyl cation mediated conversion of CO<sub>2</sub> into benzoic acid, formic acid, and methanol[J]. Angew Chem Int Ed Engl, 2012. 51(12): 2981-2984.
- [260]. Goettmann F., Thomas A., and Antonietti M. Metal-free activation of CO<sub>2</sub> by mesoporous graphitic carbon nitride[J]. Angewandte Chemie International Edition, 2007. 46(15): 2717-2720.
- [261]. Dell A. D. B., et al. Converting carbon dioxide into carbamate derivatives[J]. Chemical Reviews, 2003. 103(10): 3857-3898.
- [262]. Blanchard L. A., et al. Green processing using ionic liquids and CO<sub>2</sub>[J]. Nature, 1999. 399(6731): 28-29.
- [263]. Cadena C., et al. Why is CO<sub>2</sub> so soluble in imidazolium-based ionic liquids?[J]. Journal of the American Chemical Society, 2004. 126(16): 5300-5308.
- [264]. Li F., et al. Chemical fixation of CO<sub>2</sub> with highly efficient ZnCl<sub>2</sub>/[BMIm]Br catalyst system[J]. Tetrahedron letters, 2004. 45(45): 8307-8310.
- [265]. 张锁江等. 离子液体与绿色化学[M]. 北京: 科学出版社. 2009.
- [266]. 张锁江等. 离子液体: 从基础研究到工业应用[M]. 北京: 科学出版社. 2006.
- [267]. Blanchard L. A., Gu Z., and Brennecke J. F. High-pressure phase behavior of ionic liquid/CO<sub>2</sub> systems[J]. The Journal of Physical Chemistry B, 2001. 105(12): 2437-2444.
- [268]. Xie Y., et al. CO<sub>2</sub> cycloaddition reactions catalyzed by an ionic liquid grafted onto a highly cross-linked polymer matrix[J]. Angewandte Chemie International Edition, 2007. 46(38): 7255-7258.
- [269]. Zhang Y., et al. Cycloaddition of CO<sub>2</sub> to Epoxides Catalyzed by Carboxyl-Functionalized Imidazolium-Based Ionic Liquid Grafted onto Cross-Linked Polymer[J]. Industrial & Engineering Chemistry Research, 2012. 51: 3951-3957.
- [270]. Watile R. A., et al. Efficient synthesis of cyclic carbonate from carbon dioxide using polymer anchored diol functionalized ionic liquids as a highly active heterogeneous catalyst[J]. Catalysis Science & Technology, 2012. 2(5): 1051 – 1055.
- [271]. Tang X., et al. Polystyrene-Supported N-Heterocyclic Carbene-Silver Complexes as Robust and Efficient Catalysts for the Reaction of Carbon Dioxide and Propargylic Alcohols[J]. Advanced Synthesis & Catalysis, 2013. 355(10): 2019-2028.
- [272]. Choi H. J., Selvaraj M., and Park D. W. Catalytic performance of immobilized ionic liquid onto PEG for the cycloaddition of carbon dioxide to allyl glycidyl ether[J]. Chemical Engineering Science, 2013. 100: 242-248.
- [273]. Yang Z. Z., et al. Highly efficient conversion of carbon dioxide catalyzed by polyethylene glycol-functionalized basic ionic liquids[J]. Green Chemistry, 2012. 14(2): 519 – 527.
- [274]. Udayakumar S., Shim H. L., Park S. W., et al. Imidazolium derivatives functionalized MCM-41 for

catalytic conversion of carbon dioxide to cyclic carbonate[J]. *Catalysis Communications*, 2009. 10(5): 659-664.

[275]. Han L., Park S. W., and Park D. W. Silica grafted imidazolium-based ionic liquids: efficient heterogeneous catalysts for chemical fixation of CO<sub>2</sub> to a cyclic carbonate[J]. *Energy & Environmental Science*, 2009. 2(12): 1286 – 1292.

[276]. Shim H. L., et al. Synthesis of cyclic carbonate from allyl glycidyl ether and carbon dioxide using ionic liquid-functionalized amorphous silica[J]. *Catalysis Today*, 2009. 148(3-4): 350-354.

[277]. Gao J., et al. Preparation of polystyrene-supported Lewis acidic Fe(III) ionic liquid and its application in catalytic conversion of carbon dioxide[J]. *Tetrahedron*, 2012. 68(20): 3835-3842.

[278]. Li J., et al. Copper(II) imidazolate frameworks as highly efficient photocatalysts for reduction of CO<sub>2</sub> into methanol under visible light irradiation[J]. *Journal of Solid State Chemistry*, 2013. 203: 154-159.

[279]. Song J. L., Jiang T., Yang G. Y., et al. Synthesis of cyclic carbonates and dimethyl carbonate using CO<sub>2</sub> as a building block catalyzed by MOF-5/KI and MOF-5/KI/ K<sub>2</sub>CO<sub>3</sub>[J]. *Front. Chem. China*, 2011. 6(1): 21 – 30.

[280]. Song J. L., Hu S. Q., Wu T. B., et al. MOF-5/n-Bu<sub>4</sub>NBr: an efficient catalyst system for the synthesis of cyclic carbonates from epoxides and CO<sub>2</sub> under mild conditions[J]. *Green Chemistry*, 2009. 11: 1031 – 1036.

[281]. Miralda C. M., Zhu M. Q., Ratnasam P. et al. Zeolitic Imidazole Framework-8 Catalysts in the Conversion of CO<sub>2</sub> to Chloropropene Carbonate[J]. *ACS Catalysis*, 2012. 2(1): 180-183.

[282]. Carreon M. A. Metal organic frameworks as catalysts in the conversion of CO<sub>2</sub> to cyclic carbonates[J]. *Indian Journal of Chemistry*, 2012. 51A: 1306-1314.

[283]. Lescouet T., Chizallet C., and Farrusseng D. The Origin of the Activity of Amine-Functionalized Metal-Organic Frameworks in the Catalytic Synthesis of Cyclic Carbonates from Epoxide and CO<sub>2</sub>[J]. *Chemcatchem*, 2012. 4(11): 1725-1728.

[284]. Zalomaeva O. V., et al. Synthesis of cyclic carbonates from epoxides or olefins and CO<sub>2</sub> catalyzed by metal-organic frameworks and quaternary ammonium salts[J]. *Journal of Energy Chemistry*, 2013. 22(1): 130-135.

[285]. Talinli N, Akar A., and Aydogan A. C. A New Method for Preparation of Dihydrodinaphthopyranopyrans[J]. *Tetrahedron*, 1987. 43(15): 3505-3508.

[286]. Patel H. S. and Patel B. K. Synthesis, Characterization and Glass/Carbon Reinforcement of Acetone-Formaldehyde-1,5-Dihydroxynaphthalene-Epoxy Resin System[J]. *International Journal of Polymeric Materials*, 2010. 59(3): 151-160.

[287]. Liu Y. X., et al. Linear polystyrene anchored L-proline, new recyclable organocatalysts for the aldol reaction in the presence of water[J]. *Tetrahedron: Asymmetry*, 2007. 18(22): 2649-2656.

[288]. Liu Y. X, et al. Asymmetric Aldol Reaction Catalyzed by New Recyclable Polystyrene -supported L-proline in the Presence of Water[J]. *Catalysis Letters*, 2007. 120(3-4): 281-287.

[289]. Acevedo O. and Jorgensen W. L. Exploring Solvent Effects upon the Menshutkin Reaction Using a Polarizable Force Field[J]. *Journal of Physical Chemistry B*, 2010. 114(25): 8425-8430.

[290]. Jiang Y. L., et al. Syntheses of long-chain quaternary ammonium salts from fatty alcohols from Fatty Alcohols by Microwave Irradiation[J]. *Journal of the American Oil Chemists' Society*, 1996. 73(7): 847-850.

[291]. Ju Y. and Varma R. S. Aqueous N-alkylation of amines using alkyl halides: direct generation of tertiary amines under microwave irradiation[J]. *Green Chemistry*, 2004. 6(4): 219-221.

[292]. Romera J. L., Cid J. M., and Trabanco A. A. Potassium iodide catalysed monoalkylation of anilines under microwave irradiation[J]. *Tetrahedron letters*, 2004. 45(48): 8797-8800.

[293]. Kappe C. O. and Stadler A. *Microwaves in organic and medicinal chemistry*[M]. Wiley-VCH Verlag GMBH&Co.KGaA.. 2005.

[294]. Jastrzebski J. T., et al. The unexpected formation of a lithium-enolate from bis -[(dimethylamino) methyl] phenyl copper lithium. X-ray structure of Li<sub>4</sub>[OC(=CH<sub>2</sub>) C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>NMe<sub>2</sub>-2]<sub>4</sub> [J]. *Journal of organometallic chemistry*, 1985. 292(3): 319-324.

[295]. Pandey, et al. An improved process for the preparation of substituted acetophenones[P]. IN185806A1, 1996.

- [296].Flores K. O., et al. Microwave assisted Friedel – Crafts acylation reactions of Amberlite XAD-4&#8482; resin[J]. Materials Letters, 2007. 61(4-5): 1190-1196.
- [297].Lu X. B., He R., and Bai C.X. Synthesis of ethylene carbonate from supercritical carbon dioxide/ethylene oxide mixture in the presence of bifunctional catalyst[J]. Journal of Molecular Catalysis A: Chemical, 2002. 186(1): 1-11.
- [298].Sun J., et al. A direct synthesis of styrene carbonate from styrene with the Au/SiO<sub>2</sub>ZnBr<sub>2</sub> /Bu<sub>4</sub>NBr catalyst system[J]. Journal of Catalysis, 2005. 230(2): 398-405.
- [299].Kazarian S. G., Briscoe B. J., and Welton T. Combining ionic liquids and supercritical fluids: in situ ATR-IR study of CO<sub>2</sub> dissolved in two ionic liquids at high pressures[J]. Chemical Communications, 2000(20): 2047-2048.
- [300].Anthony J. L., Maginn E. J., and Brennecke J. F. Solubilities and thermodynamic properties of gases in the ionic liquid 1-n-butyl-3-methylimidazolium hexafluorophosphate[J]. The Journal of Physical Chemistry B, 2002. 106(29): 7315-7320.
- [301].Han L., et al. Ionic liquids containing carboxyl acid moieties grafted onto silica: Synthesis and application as heterogeneous catalysts for cycloaddition reactions of epoxide and carbon dioxide[J]. Green Chemistry, 2011. 13(4): 322 – 353.
- [302].Demberelnyamba D., Yoon S. J., and Lee H. New Epoxide Molten Salts: Key Intermediates for Designing Novel Ionic Liquids[J]. Chemistry Letters, 2004. 33(5): 560-561.
- [303].Bates E. D., et al. CO<sub>2</sub> capture by a task-specific ionic liquid[J]. Journal of the American Chemical Society, 2002. 124(6): 926-927.
- [304].Bocarsly A. B., Gibson Q. D., Morris A. J., et al. Comparative Study of Imidazole and Pyridine Catalyzed Reduction of Carbon Dioxide at Illuminated Iron Pyrite Electrodes[J]. ACS Catalysis, 2012, 2, 1684&#8722;1692.
- [305].Udayakumar S., et al. Immobilization of ionic liquid on hybrid MCM-41 system for the chemical fixation of carbon dioxide on cyclic carbonate[J]. Catalysis Communications, 2008. 9(7): 1563-1570.
- [306].Xiong Y., et al. Cycloaddition Reaction of Carbon Dioxide to Epoxides Catalyzed by Polymer-Supported Quaternary Phosphonium Salts[J]. Journal of Chemistry, 2013. 2013: 1-9.
- [307].Tan Y. X., He Y. P., and Zhang J. Serine-based homochiral nanoporous frameworks for selective CO<sub>2</sub> uptake[J]. Inorganic Chemistry, 2011. 50(22): 11527-11531.
- [308].Purohit C. S., Parvez M., and Verma S. Kinetic characterization of a bioinspired, heterogeneously active macromolecular catalyst for phenol oxidation and coupling reactions[J]. Applied Catalysis A: General, 2007. 316(1): 100-106.
- [309].Chen B. L. Metal&#8722;Organic Frameworks with Functional Pores for Recognition of Small Molecules[J]. Accounts of Chemical Research, 2010. 43(8): 1115-1124.
- [310].Perumalla S. R., Suresh E., and Pediredi V. R. Nucleobases in Molecular Recognition: Molecular Adducts of Adenine and Cytosine with COOH Functional Groups[J]. Angewandte Chemie International Edition, 2005. 117(47): 7930-7935.
- [311].Stylianou K. C., et al. CO<sub>2</sub> selectivity of a 1D microporous adenine-based metal-organic framework synthesised in water[J]. Chemical Communications, 2011. 47(12): 3389-3391.
- [312].An J., et al. Metal-adeninate vertices for the construction of an exceptionally porous metal-organic framework[J]. Nature Communications, 2012. 3: 604-609.
- [313].Verma S., Mishra A. K., and Kumar J. The Many Facets of Adenine-Coordination, Crystal Patterns, and Catalysis[J]. Accounts of Chemical Research, 2010. 43(1): 79-91.
- [314].Srivatsan S. G., Parvez M., and Verma S. Modeling of Prebiotic Catalysis with Adenylated Polymeric Templates Crystal Structure Studies and Kinetic Characterization of Template-Assisted Phosphate Ester Hydrolysis[J]. Chemistry-A European Journal, 2002. 8(22): 5184 - 5191.
- [315].Srivatsan S. G., et al. Phenol oxidation by copper-metallated 9-allyladenine-DVB polymer: reaction catalysis and polymer recycling[J]. Applied Catalysis A: General, 2001. 209: 327 – 334.

- [316].Srivatsan S. G. and Verma S. Synthetic dephosphorylation reagents: rate enhancement of phosphate monoester hydrolysis by Cu(ii)-metallated adenine nucleobase polymers[J]. Chemical Communications, 2000(6): 515-516.
- [317].Srivastava R., Srinivas D., and Ratnasamy P. CO<sub>2</sub> activation and synthesis of cyclic carbonates and alkyl/aryl carbamates over adenine-modified Ti-SBA-15 solid catalysts[J]. Journal of Catalysis, 2005. 233(1): 1-15.
- [318].An J. Design, Synthesis and Characterization of Porous Materials made from Nucleobases and Metals[D]. University of Pittsburgh. 2008.
- [319].Si X., et al. High and selective CO<sub>2</sub> uptake, H<sub>2</sub> storage and methanol sensing on the amine-decorated 12-connected MOF CAU-1[J]. Energy & Environmental Science, 2011. 4(11): 4522-4526.
- [320].Li W. J., et al. Growth mechanism and growth habit of oxide crystals[J]. Journal of Crystal Growth, 1999. 203: 186-196.
- [321].Dokocil E. J., et al. UV-Vis Spectroscopy of Iodine Adsorbed on Alkali-Metal-Modified Zeolite Catalysts for Addition of Carbon Dioxide to Ethylene Oxide[J]. Journal of Physical Chemistry B, 1999. 103: 6277-6282.
- [322].Cao H., et al. Synthesis of adenine-modified reduced graphene oxide nanosheets[J]. Inorganic Chemistry, 2012. 51(5): 2954-2960.
- [323].Balodis E., et al. Ionization constants and thermal stabilities of uracil and adenine under hydrothermal conditions as measured by in situ UV – visible spectroscopy[J]. Geochimica Cosmochimica Acta, 2012. 93: 182 – 204.
- [324].Iizumi S., et al. First observation of infrared and UV – visible absorption spectra of adenine radical in low-temperature argon matrices[J]. Journal of Molecular Structure, 2012. 1025: 43-47.
- [325].Chen Y. F., Babarao R., Li J., and Jiang J. W.. A Highly Hydrophobic Metal-Organic Framework Zn(BDC)(TED)0.5 for Adsorption and Separation of CH<sub>3</sub>OH/H<sub>2</sub>O and CO<sub>2</sub>/CH<sub>4</sub>: An Integrated Experimental and Simulation Study[J]. Journal of Physical Chemistry C, 2010. 114(14): 6602 – 6609.
- [326].Nalaparaju A., and Jiang J. W.. Molecular Understanding for the Adsorption of Water and Alcohols in Hydrophilic and Hydrophobic Zeolitic Metal-Organic Frameworks[J]. Journal of Physical Chemistry C, 2010. 114(26): 11542 – 11550.
- [327].Jiang A. N. et al. Recovery of dimethyl sulfoxide from aqueous solutions by highly selective adsorption in hydrophobic metal-organic frameworks[J]. Langmuir, 2012. 28(43): 15305-15312.
- [328].Zhan Y., et al. Silica cages with controllable frameworks: synthesis, structure-tailoring, and formation mechanism[J]. Journal of Materials Chemistry, 2008. 18(48): 5967.
- [329].Zhang Q., et al. Catalyzed activation of CO<sub>2</sub> by a Lewis-base site in W – Cu – BTC hybrid metal organic frameworks[J]. Chemical Science, 2012. 3(9): 2708 – 2715.
- [330].Wang F., et al. Hybrid zeolitic imidazolate frameworks with catalytically active TO4 building blocks[J]. Angewandte Chemie International Edition, 2011. 50(2): 450-453.
- [331].Su & kowski W. W., et al. Preparation and properties of flocculants derived from polystyrene waste[J]. Polymer Degradation and Stability, 2005. 90(2): 272-280.
- [332].Vieira G. C., et al. Sulfonated polystyrene: A catalyst with acid and superabsorbent properties for the esterification of fatty acids[J]. Fuel, 2010. 89(1): 257-259.
- [333].Bekri-Abbes I, Bayoudh S., and Baklouti M. A technique for purifying wastewater with polymeric flocculant produced from waste plastic[J]. Desalination, 2007. 204(1-3): 198-203.
- [334].Inagaki Y., et al. Reclamation of Waste Polystyrene by Sulfonation[J]. Langmuir, 1999. 15: 4171-4175.
- [335].Lee J. Y., Pan L., Emge T. J., and Li J. Microporous Metal – Organic Frameworks with High Gas Sorption and Separation Capacity[J]. Advanced Functional Materials, 2007. 17(8): 1255-1262.
- [336].Jiang H. L., Akita T., Haruta M., Sakurai H., and Xu Q. Au@ZIF-8: CO Oxidation over Gold Nanoparticles Deposited to Metal-Organic Framework[J]. Journal of the American Chemical Society, 2009. 131: 11302 – 11303.
- [337].Chen Z. J, Li M. R., Yang Q. H., and Li C.. Enhancement of the performance of a platinum nanocatalyst



confined within carbon nanotubes for asymmetric hydrogenation[J]. *Angewandte Chemie International Edition*, 2011. 50(21): 4913-4917.

[338].PAN X. L., et al. Enhanced ethanol production inside carbon-nanotube reactors containing catalytic particles[J]. *Nature Materials*, 2007. 6(7): 507-511.

[339].冉瑞成,裴伟伟,贾欣茹,沈吉. 高分子载体Lewis酸催化剂: 聚苯乙烯-四氯化锆复合物——制备及其在有机合成中的应用[J]. *分子催化*, 1988. 2(2): 112-118.

[340].Yang W., and Antonietti M. Efficient Metal-Free Oxygen Reduction in Alkaline Medium on High-Surface-Area Mesoporous Nitrogen-Doped Carbons Made from Ionic Liquids and Nucleobases[J]. *Journal of the American Chemical Society*, 2011. 133: 206 – 209.

[341].Wei J, et al. A Controllable Synthesis of Rich Nitrogen-Doped Ordered Mesoporous Carbon for CO<sub>2</sub> Capture and Supercapacitors[J]. *Advanced Functional Materials*, 2013. 23(18): 2322-2328

Degree papers are in the "[Xiamen University Electronic Theses and Dissertations Database](#)". Full texts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to [etd@xmu.edu.cn](mailto:etd@xmu.edu.cn) for delivery details.

厦门大学博硕士论文摘要库